

# Overview of SEFSC Bycatch Reduction Research

Southeast Fisheries Science Center Mississippi Laboratories Harvesting Systems Unit





# **SEFSC Gear Research Guidance**

SEFSC Sea Turtle Strategic Plan

- Development of bycatch mitigation technologies to reduce potential protected species population impacts from fishery interactions
- Develop and refining modifications to fishing gear and practices that can reduce sea turtle bycatch without implementing fishery closures
- Inform amendments to fishery regulations pertaining to modified gear or practices and train enforcement personnel
- Conduct training to ensure proper use of gear by fishers




# **SEFSC Gear Research Guidance**


- Magnuson-Stevens Fishery Conservation and Management Act;
   National Standard 9; Bycatch
- Marine Turtle Conservation Act (2004)
- NMFS Strategic Plan, SEFSC Strategic Plan
- 2014 Biological Opinion Southeast U.S. Shrimp Fisheries
- NMFS Bycatch Reduction Engineering Program
  - Participate as SEFSC representative
  - Develop national priorities for bycatch reduction research
  - Direct funding to internal (NMFS) and external bycatch reduction researchers



# SEFSC Harvesting Systems Unit Background

- Unique mission within NOAA Fisheries
- Gulf of Mexico, SE Atlantic & International
- 4 Research fisheries biologists
  - 3 Gear specialists
- Fisheries: Trawl, Longline, Gillnet
- Specialized techniques
  - SCUBA
  - Autonomous U.W. cameras
  - High resolution sonar







# **SEFSC Gear Research**

Fishery	Species	Research
Trawl (shrimp, flounder, flynet)	turtles	TEDs
Gillnet	turtles / sturgeon	Low profile, lights, modified panels
Longline	turtles	Circle hooks, bait type, soak time
Trawls (shrimp)	mammals	Behavior assessment



### **Research Methods**

# Fishery-Dependent Research

- At-sea testing and evaluation aboard commercial vessels
- Assess usability of new gear concepts
- Evaluate effects on target catch



# Fishery-independent research

- Proof of concept evaluations aboard NMFS R/Vs
- Using advanced technology to:
  - Evaluate gear dynamics
  - Understand fish and marine animal behavior relative to gear





# Approving new TEDs: Juvenile turtle testing protocol

Controlled test using NOAA divers and captive-reared, 2-yr old loggerheads

Candidate TEDs are certified through the protocol, which was published and has been in use since 1990 (FR Vol. 55, No. 195)

Provides opportunity for industry and researchers to evaluate new ideas



Resulted in significant improvements in TED performance with regard to turtle exclusion and shrimp retention

# **Recent Research: Skimmer Trawl TEDs**

### **Small Turtle Exclusion Rate Evaluations**

Captive-reared, one year old, loggerhead sea turtles are used as surrogates for Kemp's ridleys caught in the fishery to determine exclusion rates for various TED configurations.



### **Independent Catch Retention Testing**

Suitable TED configurations identified through small turtle evaluations are tested under commercial conditions aboard NOAA vessel *RV Caretta* to determine levels of shrimp loss and bycatch reduction associated with each.



### **Dependent Catch Retention Testing**

TED configurations that exhibit minimal shrimp loss during independent testing are further tested under various commercial conditions aboard multiple commercial vessels throughout the southeastern skimmer trawl fishery.



# Flynet TED Research

# **Challenges Transferring TED Technology**

# **High Catch Rates**

Flynets are high opening bottom trawls utilized to target various species of finfish including Atlantic croaker. Catch rates may exceed 20,000 lbs in 30 minutes.

# **Gear Storage on Net Reels**

Nets and associated gear are deployed, retrieved and stored on stern mounted net reels, which does not lend to the use of rigid frame TEDs.

# **Handling and Long-Term Durability of Gear**

Large catches combined with rough seas require TED designs that are virtually "hands-free" and extremely durable.







# Flynet TED Research

# **TED Design and Development**

### **Design development**

Multiple prototypes evaluated with research focused on flexible designs.









### **Three Pronged Approach**

Prototype designs are evaluated for:

Turtle Exclusion – Small turtle testing certification

Usability – Installation and use in standard commercial gear

Catch Retention – Trouser trawl or alternate haul





# **Certified Designs**

Flexible Flatbar Flynet (FFF) TED – Contains a flexible center section with target catch reduced by 3.9%


**Hopkins Cable TED** – Constructed completely of cable, scheduled for catch retention testing fall 2015.




# Overview of Experimental Gillnet Research – 2001 – 2014

### **Modifications Assessed:**

- Low Profile Gillnets (NC, Trinidad)
- Double Lead line Gillnets (NC)
- Modified Panel Gillnets (VA)
- Lighted Gillnets (NC)



# Purpose/Methods




Tested modified gillnets to reduce protected species interactions

Species of concern included sea turtles, sturgeon and other finfish bycatch

Experiments conducted in NC, VA, and Trinidad in estuarine and nearshore waters

Matched paired sets between control and experimental nets

Studies were conducted both independently and dependently



# Results

Low profile gillnets proved effective (38%, 80% reductions in sea turtles)in Trinidad and NC

Modified panels (3.5' openings) reduced sturgeon by 64%, and increased striped bass 46%

Double lead line and gillnet light studies were inconclusive due to area tested in NC


# **Key Points / Future Research**

Tie-downs in gillnets increase entanglement and potential mortality

If target catches can be maintained, less webbing in the water column is desirable

No one-size fits all net configuration; varies by depth, fishery, location


Continued research with lighted gillnets, rectangular mesh, and double lead line needed









# Sea Turtle Bycatch Experiment Northeast Distant Waters Swordfish Fishery 2001-2003





# Pelagic Longline Sea Turtle Bycatch







Leatherback turtles generally entangled or foul hooked




# **Bait Type:**

Fish vs. Squid


# **Hook Type:**

J-hook vs. circle hook

# **Sea Turtle Bycatch Experiment NED Swordfish Fishery 2001-2003**









# **Optimal Combination For Sea Turtle Mitigation**







18/0 Circle Hook

# Sea Turtle Bycatch Experiment Results

	% Difference
Loggerhead Turtles	- 88% (reduction)
Leatherback Turtles	- 63% (reduction)
Swordfish	+ 20% (increase)
Bigeye Tuna	- 80% (reduction)

Watson. et al. 2005. Fishing methods to reduce sea turtle mortality associated with pelagic longlines. Can. J. Fish. Aquat. Sci. 62: 965-981.



# Weak Hook Experiment Results


- 8 commercial vessels involved in tests
- 418 pelagic longline sets completed
- 134 bluefin were caught during the experiment
- 47 caught on the experimental hook
- 46% reduction in bluefin with weak hooks
- 2,547 yellowfin tuna landed
- 6% reduction in yellowfin catch (ns)
- Majority of bluefin escapes took place in < 5 min.</li>







# Reducing bycatch in the reef fish bottom longline fishery





# **Elapsed Time (Hook Timer)**








# **Bite Times**

Reef Fish vs. Sharks



Shorter soak time reduces bycatch

#### GROUP

- RED GROUPER
- RED SWAPPER
- SHARK



# **NEFSC Gear Research**

Fishery	Species	Research
Trawls (Flounder & Scallop)	turtles	TEDs, Topless trawls, data loggers
Gillnet	sturgeon	Low profile and stand-up gillnets
Gillnet	turtles	Low profile gillnets
Scallop dredge	turtles	Chain mats and deflector dredge
Pound net	turtles	Modified leader
Longline	cusk	assessment of barotrauma



# Collaboration with NEFSC Gear Research Program: Scallop Fisheries

Assisted NEFSC with initial evaluations of industry developed scallop dredge and modified TEDs to reduce turtle bycatch







# Collaboration with NEFSC Gear Research Program: Flounder TEDs



23/05/2007

- Participated in flume tank testing
- Assisted in rigging and training of TED installation
- Conducted small turtle certification for flounder TEDs




# **Future collaborations with NEFSC**

- Fall/Winter 2015
- Fishery-dependent
- Flexible cable TEDs
- Mid-Atlantic croaker and flounder testing for catch retention







# WET LAB FACILITIES Galveston Lab




Raceways & leatherback filtration system





Water treatment
Includes mechanical
& biological filters
+ protein skimming
with ozone and U/V
Sterilization





Water chiller units





Turtle buildings





Flow-through seawater basins with 2 year-old loggerheads

# PANAMA CITY LAB FACILITIES





Sea turtle "conditioning pens"



# **SEFSC Gear Research**

### Successes

### **TEDs** – shrimp fishery

- 97% effective in reducing turtle captures in shrimp trawls
- No effect on target catch rates with optimal TEDs
- International adoption of technology

### **TEDs** – flounder fishery

Specialized TED minimized catch loss and 97% turtle exclusion

#### **Circle Hooks**

- Significant reductions in turtle interactions in NED fishery
- Fishery closure prevented
- Increased swordfish catch rates



# SEFSC Gear Research

### **Limitations**

### **Permitting**

Limits fishery-independent directed research

#### **Commercial vessel charters**

- Contracting process overwhelming for owner/operators
- 1-yr lead time for contracts: difficult with temp. funds

### **Industry participation**

- Controversial issues can effect active involvement
- Researchers act as agency liaisons on bycatch issues
- Improved communication: agency management to industry



# **SEFSC Gear Research**

# **Recommendations and Improvements**

### Early and sustained communication and planning

SERO – Researchers – Industry – State agencies

#### **Enhanced research collaborations**

- International researchers
- Expansion of NMFS BREP: promote internal collaborations

#### Fiscal re-allocations

- Funding streams often conflict with fishing seasons
- Increased ability to carry over research funds needed